

Effect of Serum Trace Elements in Hand Eczema Patients in Sulaimani Province: A Case-Control Study

Nasrin Sulaiman Hamad¹ and Diary IbrahimTofiq²

¹Dental Nursing Department Sulaimani Polytechnic University- Technical Institute of Sulaimani, Slemani, Kurdistan Region, Iraq

²Chemistry Department, College of Science, University of Sulaimani, Slemani, Kurdistan Region, Iraq

Abstract. Atopic dermatitis (AD) pathophysiology incorporates a complex interaction exchange of brokenness of the immune system, science, and environmental factors. It is notable that nutritional status is significant for an appropriately working immune system, which prompts a furiously contested inquiry regarding the part of dietary variables in the pathogenesis of AD.

Minor components are micronutrients found in the body in restricted sums, and are a vital factor in keeping up a healthy immune system. In this investigation, we examined the degrees of to the (Mg, Ti, Mn, Fe, Zn, Cd, Cr, Cu, Ni, Pb, Al, and Se) in the serum of the AD understanding we contrasted our discoveries and those of a healthy group.

The investigation populace included 60 AD patients and 50 controls from the center of skin feeding in Slemani, private skin clinic, and clinical center of many concrete production lines. We performed minor components estimation in the serum tests acquired from the two patients and the control group utilizing (ICP-MS).

We found a significant difference ($P < 0.05$) of Mg, Mn, Fe, Zn, Cd, Cr, Cu, Ni, Pb, and Se in the serum levels of the AD patients in comparison to the control group, and showed a reduction difference for the Mg, Ti, Mn, Fe, Zn, and Cd (35.576 ± 4.67 , 299.989 ± 59.64 , 133.471 ± 80.17 , 3316.234 ± 1132.78 , 289.507 ± 61.23 , 28.304 ± 35.50 , and 18.509 ± 6.703) correspondingly, while for the Al, Cr, Ni, Cu, Se and Pb (2599.950 ± 559.57 , 45.342 ± 17.69 , 463.838 ± 626.69 , 1524.524 ± 547.82 , 31.775 ± 18.11 and 180.717 ± 127.47) respectively, revealed significant increasing.

Assessment of the zinc and lead serum levels in the dermatitis populace might be valuable, and further examination of their immunological capacity in pathogenesis and treatment of skin inflammation is required.

Introduction

Atopic dermatitis (AD) is an inflammatory skin infection that includes complex collaborations between immunological, genetic, and ecological impacts. The commonness of AD has expanded internationally throughout the most recent 30 years, influencing 20% of children and 3 percent of grown-ups around the world [1]. It is firmly connected to asthma and unfavorably susceptible sharpening, youngsters who originate from families with an AD history, asthma or hay fever are bound to create skin inflammation [2]. Proper nutrition is an important factor in immune health especially the sufficient intake of vitamins, minerals and components [3]. For a long time, the association among sustenance and AD pathogenesis has been discussed [4].

Minor components are synthetics that in living beings and biological systems are available in various structures in a characteristic material at low concentrations not exactly ($\mu\text{g/g}$). In science, a minor component is a dietary mineral needed in exact moment amounts for the creature's appropriate development, advancement and physiology [5]. Minor components are evaluated as important or insignificant as per whether a life form may develop in its nonappearance and complete its life cycle [6]. Minerals and essential components are healthfully fundamental segments. These components assume a critical part in the typical working of the immune system and antioxidant systems related with atopic dermatitis pathogenesis [7]. They practice their capacity through association with biomolecules in the human body [8]. A wide assortment of minor elements are fundamental for entire body cell measures, including cell turnover, cell digestion, and apoptosis. A few examinations contemplated the linkages among AD and different minerals, including zinc, copper, iron and magnesium [9].

There stays a conclusive connection between zinc lack and AD, which seems to assume a part in the infection cycle [10]. Studies on the iron status of AD patients indicated lower serum iron levels than those of control groups [11]. There are few studies of Mn Allergy patients [12]. Concerning hypersensitive chromium-caused contact dermatitis, distributed examinations have appeared over all sure relationship with atopic dermatitis [13]. A couple of studies have arisen on the connection between intrauterine presentation to lead and atopic sickness advancement in earliest stages. Nonetheless, as far as anyone is concerned, no investigation has yet been directed on a potential relationship between intrauterine cadmium presentation and the advancement of atopic skin inflammation in earliest stages [14]. Animal models detailed that mice took care of a magnesium-inadequate eating routine had expanded degrees of pro-inflammatory cytokines and skin sores created with an example like AD [15]. This examination meant to dissect the groupings of minor components in serum including; Mg, Ti, Mn, Fe, Zn, Cd, Cr, Cu, Ni, Pb, Al, and Se and to assess their significance to infection severity in a grown-up with AD and furthermore contrasted with the healthy control group.

Patients and Methods

The study population consisted of 60 patients with hand eczema (22 females and 38 males aged 17-62 years) at various workplaces (27 from the Taslwja and Gasin cement plants in Slemani, considered to be industrial areas, and the rest from the non-industrial zones). Fifty safe participants representing the control group (25 men with 25 women aged 20-61 years) were also included in the study; individuals in this group were free of any signs and symptoms of atopic dermatitis or no serious medical disorder or treatment over a 3-month span at the time of blood collection. The respondent admitted to the Slemani Governorate Skin Feeding Center.

Experimental design

In order to ascertain the presence of any diseases that could influence the parameters of the test, multiple test patient clinical trials were performed. For all sample populations, regular physical examinations, including organ function, dietary status, weight, blood pressure, and chest X-rays were performed, and special data (Questionnaires) were collected from information prepared for these purposes from registered persons (Table 1). The research populations were classified into five major age groups: Group A (age 15-25 years); Group B (age 26-35 years); Group C (age 36-45 years); Group D (age 46-55 years); and Group E (age 65 years).

Using the atopic dermatitis classification system (SCORAD), the patient was assessed for severity. AD is ranked mild (SCORAD <25), moderate (SCORAD 25-50) or severe (SCORAD over 50) by SCORAD.[16]. A stable individual was part of the control group. This research was approved by the local ethics committee.

Table 1: The questionnaire employed in the sampling campaign.

1- Subject numbers
2- Name, and Gender
3- Smoking history
4- Level of education
5- Residency: Industrial (If the industrial time of employment) or not industrial
6- Alcoholism
7- Vegetarian
8- Weight
9- Height
10- Taken VD before
11- Family history
12- Duration of illness
13- Chronic diseases:- Hypertension, cholesterol, diabetes, heart disease, asthma, hypothyroidism, obesity, and chronic kidney disease.

Blood sample collection and serum separation

Venous blood (5 ml) was acquired in an ethylenediaminetetraacetate tube and drawn from every member. At 3000 rpm the centrifugation was directed for 10 min to detach the serum. Until quantitative

examination the serum was put away at -20°C . Agreement with the arrangements to prerequisites set by the National Committee for Clinical Laboratory Standards, steps were taken to eliminate any metal defilement during blood assortment, handling and capacity.

Determination of serum trace elements level

The ICP-MS estimations were finished utilizing an (Agilent 7500 ICP-MS, furnished with ASX-520 Autosampler) in Zaminrizkavan Co. Ltd. Focus in Taran, Iran. Tests (0.2 ml (or 200 μL) were weakened with deionized water and dilution component (A dilution variable for 50x considered in the subsequent information, so no back-estimation to the concentration of the investigation in the first examples required), and a volume of 10 ml utilizing supra unadulterated nitric acid (0.5%), which is set up from 99.9% supra unadulterated nitric acid. At that point every serum test experienced a 0.45 μm channel to eliminate any suspended material prior to running examples.

The alignment was performed utilizing a clear and six standard focuses (50, 20, 10, 5, 2, and 1 mg/L) from a combination of 2 "ICP-OES/MS multi-component. Absorbance was estimated at 213.8, 327.5, 279.9, 421.5, and 286.2 nm, and The accompanying components were estimated with the ICP-MS: Mg, Ti, Mn, Fe, Zn, Cd, Cr, Cu, Ni, Pb, Al, and Se individually. "0" means "clear example" (weakened supra unadulterated nitric corrosive) without arrangement standard components as appeared in figure 1.




Figure 1: Calibration curves

Statistical analysis

Information were analyzed utilizing the Statistical Package for Social Sciences (SPSS, variant 22). The T-test was utilized to analyze the methods for all boundaries, minor component concentrations in the various groups (controls and dermatitis patients) in the investigation. Moreover utilizing statistical regression methods for clarifying the connections between totally considered minor components with all boundaries explicitly the most significant elements (Zinc and Lead). A p-estimation of ≤ 0.05 was considered measurably huge.

Results

Population characteristics

A total of 110 participants were included in the study. Sixty of them showed all symptom of eczema that composed of 22 females (36.7%) and 38 males (63.3%), in different location of work 27 (45%) worked in the industrial zone and 33 (55%) in the non-industrial zone, while fifty out of 110 were in healthy condition that composed of 25 males and 25 females (50%), 7 (14%) of them are from the industrial zone and 43 (86%) from the non-industrial zone, with five groups of ages arranged from less than 20 to more than 51 years old.

Table 2 utilizing the SCORAD record, 19 cases (31.7%) with AD were classified as having a mild AD with the value of (24.23±7.01), 23 people (38.3%) were arranged as having a moderate disease with the value of (17.54±2.05), and 18 people (30.0%) were ordered as having marked AD with the value of (12.40±1.58).

Table 2: Frequency and severity of AD cases.

AD	SCORAD index		P-value
	Number %	Mean ±SD	
Mild	(19) 31.7%	24.23±7.01	0.000
Moderate	(23) 38.3%	17.54±2.05	
Severe	(18) 30.0%	12.40±1.58	

The association effect of some parameters between control groups and eczema patients

A comparison of various parameters among the population participated in this study (control groups and eczema patients) were shown in Table 3. The statistically significant difference between the means of each parameter (education level, place of work, and family history) was noted by the (P= 0.000, 0.000, and 0.013) respectively, in control and patient groups. While there was no statistically significant difference between the means of the other parameters between healthy and AD groups.

Table 3: The association effect of some parameters between control groupseczema patients by using (T-test).

Parameters	Mean± S.D		T-value	P-value
	<i>Control</i>	<i>Eczema</i>		
Age	36.08±9.40	36.95±11.91	0.42	0.676
Smoker	0.34±0.48	0.35±0.61	0.10	0.923
Level of education	2.48±0.76	1.45±1.00	-6.13	0.000
Location of work	0.14±0.35	0.46±0.50	3.87	0.000
Alcoholism	0.18±0.39	0.07±0.25	-1.85	0.068
Vegetarian	0.04±0.20	0.03±0.18	-0.18	0.854
Chronic disease	0.78±2.44	231.80±1600.26	1.12	0.268
Family history	0.00±0.00	0.10±0.30	2.56	0.013

The results are expressed as mean±S.D, (No. patient =60 and No. control =50), $P \leq 0.01$.

Influence the relationship of serum trace elements on the prevalence of eczema disorders in comparison to the control group

Patients with AD had a significantly moderate reduction in serum Mg, Ti, and Mn elements levels (35.576±4.6, 299.989±59.64 and 133.471±80.17) consecutively, in comparison to the healthy individuals (39.669±8.42, 372.546±73.07 and 167.491±437.84) as seen in figure 2A-C. While, the serum concentration of Fe, Zn and Cd significantly were lower in AD cases as shown in figure 2D-F (3316.234±1132.78, 289.507±61.23 and 28.304±35.50) successively, in contrast to the normal population group (3499.085±1470.98, 725.069±181.96 and 77.842±39.54). Significant increasing of Al, Cr, and Ni serum levels(2599.950±559.57, 45.342±17.69 and 463.838±626.69) were found in the eczema patient in comparison to the in the control group(1980.147±426.87, 20.817±4.92 and 28.961±7.67) as seen in figure

2G-I. Finally, a statistically significant increase in the level of serum Cu, Se and Pb (1524.524 ± 547.82 , 31.775 ± 18.11 and 180.717 ± 127.47) consecutively, were observed in AD patients when compared with the control group (1118.868 ± 186.46 , 0.799 ± 1.09 and 88.067 ± 26.05) as revealed in figure 2J-L.




Figure 2: The column chart showed serum trace elements level on the prevalence of eczema patients in contrast to the control group.

Regression analysis

In this study regression analysis used to identify the strength effect of some parameters such as Gender, level of education, location of work, family history, BMI, and severity (independent variables) on the concentration of all elements found in table 4.

Hand eczema tended to increase significantly in gender for each (Mg, Al, Ti, Cr, Fe, Ni, and Cd) minerals with exception of (Mn, and Se) revealed that were no statistically significant differences. The prevalence of atopic dermatitis increased significantly regarding the level of education in the following minerals; Al, Ti, and Cu. While the other elements showed a non-significant association. The T-test and p-value for each element according to residency revealed that (Cu, Cd, and Se) were not statistically significant. While the prevalence of hand eczema among other metals presented the statistically significant differences between a person who was work in different industries and exposed to the minerals including; (Mg, Al, Ti, Cr, Mn, Fe, Ni, and Cu). Concerning the family history among all the elements, only Ni showed a highly significant difference, whereas the remainder showed a non-significant difference. Conferring to the severity of patient only Mg and Se amongst all the metals exhibited a highly significant difference, however, the other metals showed no statistically different between the severity of the patient.

Table 4: Regression results of the effectiveness of different parameters on the various trace elements concentration in eczema patients.

Variables	R-square	T-test		F-test		Beta
		T-test	P-value	F-test	P-value	
		8.335	0.000			.618
		1.795	0.078			.099
Gender	0.94	3.922	0.000	179.006	0.000	.180
Level of education		Mg	-.267			0.790
Residency		2.132	0.037			.184
Family history		7.157	0.000			.612
Severity of patient		2.907	0.005			.184
	0.92	3.167	0.003	132.287	0.000	.167
		.442	0.660			.019
		1.033	0.306			.103
		6.878	0.000			.600

			2.958	0.005			.192
	Ti	0.9	2.319	0.024	126.420	0.000	.125
			.206	0.837			.009
			1.367	0.177			.139
			4.412	0.000			.540
	Cr	0.84	1.238	0.221			.112
			2.021	0.048	58.830	0.000	.153
			-1.093	0.279			-.066
			1.632	0.108			.232
			1.837	0.072			.287
			1.883	0.065			.219
	Mn	0.74	2.269	0.027	31.780	0.000	.219
			.543	0.589			.042
			1.338	0.186			.243
			4.221	0.000			.517
	Fe	0.84	.809	0.422			.074
			2.300	0.025	58.670	0.000	.174
			.300	0.766			.018
			1.733	0.089			.247
			-2.407	0.020			-.366
			1.310	0.196			.155
	Ni	0.29	-3.196	0.002	4.460	0.002	-.485
			2.320	0.024			.284
			.171	0.865			.020
			8.414	0.000			.819
	Cu	0.9	2.777	0.007			.200
			2.079	0.042	99.540	0.000	.125
			1.076	0.287			.052

Table 5: Regression analysis of the effectiveness of different parameters on the Zn and Pb concentration in AD patient.

Parameters	R-square Zn and Pb	T-test		F-test		Beta
		T-test	P-value	F-test	P-value	
Gender	0.74	17.78	0.000	315.97	0.000	0.862
	0.59	12.73	0.000	161.96	0.000	0.77
level of education	0.75	18.27	0.000	333.85	0.000	0.868
	0.48	9.95	0.000	99.0	0.000	0.690
Location of work	0.34	3.82	0.000	14.62	0.000	0.344
	0.43	5.03	0.000	25.33	0.000	0.434
Family history	0.19	-1.93	0.046	3.72	0.046	-0.182
	0.38	4.31	0.000	18.55	0.000	0.381
BMI	0.75	18.01	0.000	324.37	0.000	0.865
	0.63	13.54	0.000	183.37	0.000	0.792
Severity of patient	0.84	1726	0.000	397.95	0.000	0.914
	0.48	7.44	0.000	55.33	0.000	0.696

Analyzed by regression test, * $p < 0.01$. The first showed all values for Zn, while the second row revealed values for the Pb.

Discussion

Investigation into the part of micronutrients on AD has a few constraints. Over late years interest has expanded in the effect of nutrition on AD and in the function of micronutrients specifically [17]. The interchange between the immune system, cell capacity, vitamins and minerals is unpredictable; it is hard to close the disconnected impact of a solitary or scarcely any mixes[9].

This case-control study gives the principal proof to the relationship between serum minor components and AD in the Kurdish populace. As per this investigation, the female is less danger to dermatitis contrasted with guys, And errors with the past examination recommended that a high commonness of AD in females than guys might be because of the maltreatment by females of makeup as they are more worried about their skin appearance than guys[18], Other examination have exhibited that skin inflammation is twice as

far reaching in females as in guys[19]. The motivation behind why the event of females diminished in this investigation could be because of a restricted information populace or introduction of guys to substance material than females because of their working style or occupation, which adds to dermatitis creation

Air contamination is the reason for a wide scope of modern and non-substance poisons. Air toxins can begin inside and outside and can infiltrate the skin, tie to the layer corneum and enter the fundamental flow [20]. On the side of our discoveries, the predominance rates in created countries have ascended to as high as 15-30 percent of youngsters and 2-10 percent of grown-ups in the course of recent many years[21]. The past examination underpins our discoveries, where there was a solid connection between the working environment and the event of dermatitis, an aggregate of 32 out of 80 laborers (40%) at the Bank of Greenland created skin issues when moving to redesigned premises, 22 out of 27 had improved at first whining of skin inflammation, and all instances of hand dermatitis had been cleared. The beginning of manifestations of building redesign, the particular objections and the vanishing of side effects after work environment mediation unequivocally show a connection between remodeled assembling presentations and dermatitis[22].

Concurrence with our outcomes, the past examination detailed that the greater part of the investigation bunch had optional and advanced education, proposing a more noteworthy arrangement and worry about atopic dermatitis [19]. Contrasted with our past examination with a set number of patients, schooling in youngsters with atopic dermatitis and their folks has had a valuable impact [23, 24].

It is perceived that the commonness of AD in urban communities is higher in populaces of a similar nationality and hereditary history than in the countryside. Applicable ecological danger factors are urbanization, aberrations in hygiene, microbial diseases, vaccination, anti-toxin utilization, environmental contamination, sensitivity to allergens and diet [25]. Following our outcomes, no doubt toxin introduction here and there secures against the advancement of AD or that presentation to exposure to infectious diseases related with poorer, less developed nations protects against AD [26]. Skin inflammation is additionally more common in metropolitan versus provincial networks and seems to target offspring of a higher financial status in more modest families[27, 28].

Consent to our examination, past information revealed that there was a strong relationship among parental and youth AD, paying little heed to which parent had AD [29]. Our outcomes uphold the hypothesis that dermatitis has a polygenic etiology and show that qualities related with parental skin inflammation are all the more firmly identified with kid skin inflammation [30]. A strong family background of hypersensitive illnesses (47.5 percent) was found in the investigation among AD patients [31]. Family ancestry is one of the primary symptomatic models for AD, and it is viewed as a critical danger factor for AD. Studies from the West discovered a sound family background of AD in 53-80 percent of patients[32, 33]. In a grown-up beginning AD study, 54.2 percent had a family ancestry with most instances of hypersensitive rhinitis[34].

Magnesium is a fundamental minor component which assumes a significant part in the immune reaction. Magnesium is a fundamental minor component which assumes a significant function in the immune reaction.[35]. In concurrence with the current examination, past reports archived that the serum magnesium levels in patients were lower than in the benchmark group, and magnesium insufficiency was more successive in the patient group[11]. Animal models uncovered that mice had expanded degrees of supportive of fiery cytokines took care of with a magnesium-insufficient eating regimen and created skin

injuries like those of AD[15]. A difference to the discoveries acquired in this investigation, Ertunç et al. (2003) explored serum magnesium levels in kids and grown-ups with AD and found no contrasts among patients and controls [36]. Tuchinda et al. (2014), once in the past, reported that immediate contact with Mn was essentially identified with AD advancement[37]. As opposed to our investigation, Kim et al. (2017) demonstrated that the degree of Mn in the hair test of dermatitis quiet was factually higher contrasted with the control group, while different investigations showed that there was no distinction in the degree of Mn in hair tests between the two groups of skin inflammation patients and solid or relationship with clinical severity[38].

Also, in the current examination, Fe, Zn, and Cd fixation were diminished fundamentally in their level in dermatitis cases rather than the healthy populace groups as uncovered in Figure 2D-F. As per our information, comparative reports demonstrated that the absolute mean estimation of Fe and Zn in the scalp tests of dermatitis patients was decreased, this could be identified with the way that skin illness could be brought about by iron inadequacy, just as that iron might be lost in the quickened keratinocyte turnover from scaling; as of now, iron malabsorption is acknowledged as representing dermatopathicanemia[18, 39]. In contrast with our discoveries, Kaur et al. (2001) demonstrated an expanded cell turnover in AD skin, in which iron creation may increment and iron focuses in AD dermis may increment[40]. Supporting for our information, concentrates on the iron status of patients with AD have demonstrated lower serum iron levels than the populace healthy [1, 11]. Numerous examinations explore the connection between Zn levels and atopic dermatitis. In another examination on atopic dermatitis, it was discovered that erythrocyte Zn levels, which were professed to be a superior proportion of mellow Zn lack, were essentially lower in patients with atopic dermatitis contrasted and those of healthy control subjects[11]. Zinc lacks are very basic in individuals living in helpless nations who don't have adequate supplements and nutrients to keep up great wellbeing [41]. As opposed to our outcomes, Cd in AD patients was likewise essentially higher than in control patients in the Kim et al. (2017) study[38], also, different past investigations didn't show any connection among's cadmium and AD movement[42].

In this investigation, Al, Cr, and Ni components demonstrated notably rose rather than the benchmark group as found in Figure 2G-I. Past information upheld our examination by expanding the Al level in skin inflammation patients in contrast with the healthy group[38]. Components recently answered to cause AD incorporate Cr and Ni, these components are haptens that expect clinging to protein atoms to frame antigenic edifices to empower dendritic cell acknowledgment, prompting sharpening and improvement of AD [43]. In an investigation of 137 youngsters with dermatitis, 19.3 percent of patients were positive in metal fix tests and the examination affirmed that Ni and Cr are the two most as often as possible identified primary metal hypersensitivities in the skin inflammation populace[44]. As indicated by the past pilot investigation of a progression of clinical cases, which found that the word related treatment of DMG test-positive instruments, keys and other nickel-delivering items significantly expanded the nickel substance of presented skin comparative with a non-uncovered skin control site[45, 46].

While in the current investigation uniquely expanding in their mean of Cu, Se and Pb were found in the dermatitis populace in relationship to the control cases as found in Figure 2J-L. Our information upheld by various investigations, in which the serum Cu level in extreme instances of the infection was moderately high contrasted and the control [47, 48], Certain tests have indicated more significant levels of copper in grown-ups and kids with AD than in controls, albeit different tests didn't show any relationship

with seriousness of the illness[49, 50]. The authors demonstrated that significant levels of copper could result from expanded degrees of ceruloplasmine during skin aggravation [11]. Contrasts in dietary propensities, age, and procedure of examination can result from the distinction between our discoveries and the clinical writing. A previous investigation detailed that the measure of Se hair indicated a non-huge decline in AD patients contrasted with control patients, though a similar report found that Pb was altogether higher in AD patients contrasted with control patients [38]. Nonetheless, different investigations have exhibited positive or negative relationship with Se serum fixations [51, 52]. Our perception uncovered that men have higher serum Se than ladies, in concurrence with information formerly detailed [53, 54] There is, in any case, another investigation that recorded comparative Serum Se esteems for the two people [51].

Zinc is a basic micronutrient engaged with managing the intrinsic and versatile reaction to resistant frameworks. Unhealthiness is the essential driver of the zinc insufficiency. Zinc inadequacy prompts cell-intervened immune dysfunctions, among different appearances [55]. Pb is answerable for debilitating the cells that make up the human immune system (T and B lymphocytes, macrophages, and Langerhans cells). Moreover, Pb can influence the cell as well as the humoral safe reaction by lessening immunoglobulin creation, accordingly inclining individuals to expanded inflammatory diseases[56]. In this investigation, the mean serum Zn and Pb fixation demonstrated a huge relationship among sex as appeared in table 5, which is following the past reports in the Brazilian populace [57]. Males can likewise have a higher substance of the Zn minor components because of word related presentation as they work in ranches or through open air introduction to clean. Not at all like our outcomes Kim, H-j, et al indicated no huge contrasts in the centralization of Zn among people in everyone of Korea Korea[58]. Differences in serum zinc levels by age have been reported before[59]. Maternal instruction is known to substantially affect kid endurance and healthful results, the past paper found that tutoring for ladies at a worldwide level is associated with lower zinc in everyone [60]. As to, our examination upheld by the past investigation, where additionally the Pb level was higher in female than male sex [58], While, as opposed to different examinations, it was reported that blood Pb focuses in Korean populace were higher in guys than in females[61], these varieties in Pb rates among male and female subjects might be related with contrasts in way of life and word related presentation to Pb. Studies have additionally indicated that Pb seems to gather more in guys than females, prompting higher blood levels of Pb because of word related presentation [62]. Our finding introduced critical contrasts in instruction level and family ancestry with Pb level, which is in affirming with past information, who proposed that they may be frustrating factors and had a decent connection with one another [63].

Our investigation demonstrated a huge impact between the mean serum of Zn and Pb fixation and area of work as uncovered in table 5. Mohmand, J., et al. (2015) evaluated levels of conceivably harmful minor components including; zinc, examined in provincial, metropolitan, and mechanical zones of Punjab, Pakistan, presentation in all examples indicated altogether higher centralizations of Zn minor components than those from modern (which, on the other hand, had higher convergences of Pb) than country tests. Bioaccumulation patterns uncovered that presentation to tidy was one of Pb's principle courses into the human body, though Zn 's weight could be all the more firmly identified with dietary sources. The centralizations of minor components in the Punjab tests were practically identical and additionally higher than those recorded all around the world [64]. On the side of our investigation, the past discoveries demonstrated that high Zn content in the dirt examples examined proposes that the amassing of these hefty metals in the locale got essentially from the processing plants in the area, either from the removal of

modern waste or from the arrival of unregulated effluents into the dirt defiling water bodies and streams [63]. A few writing information demonstrate that huge potential reasons for metal refinement, for example, Zn, are metal inserts utilized for different clinical purposes, for example, intravascular stents, dental inserts, heart pacemakers or gynecological inserts [65]. As per our examination, Carbal Pinto et al. (2019) expressed that the elevated level of Pb metal in the dirt was essentially because of the previous mechanical action and rural practices that altogether decreased soil quality. Debasement of soil quality by contamination of metals/metalloids causes ecological dangers, causes contamination of the groundwater, is hurtful to human wellbeing and diminishes the nature of human life [66]. The situation of working and living with improved metal hence had a high effectiveness in growing AD.

The current examination indicated likewise a profoundly huge relationship between family ancestry and the serum level of Zn as found in table 5. In opposition to our discoveries, the past forthcoming investigation found that the maternal birth companion surveyed the relationship between newborn child or maternal hair Zn mineral substance at multi month old enough and AD conclusion at 10 months old enough after acclimation to family background of AD, neither baby nor maternal hair zinc at multi month old enough was identified with AD status at 10 months old enough [67].

Our information indicated a critical connection among BMI and the serum level of Zn and Pb. The connection between zinc levels and weight file (BMI) was affirmed in the past investigation, and the serum zinc levels were lower in kin with expanded weight file [68]. No doubt raising the Zn status may effectsly affect controlling food consumption [69]. One of the instruments recommended might be connected to the gainful effect of raising the Zn status on leptin control to hinder eating practices by diminishing the measure of neuropeptide YmRNA[70]. Moreover, our examination supported past investigations, in which an elevated level of Pb was found in people with expanded weight record paying little heed to age [71, 72], While our outcomes were not in understanding, there was a negative relationship among lead and stoutness or overweight populace[73].

Moreover, Zn and Pb levels indicated a solid relationship in our outcomes as appeared in table 5 in struggle with past reports that demonstrated frail connections with long haul sickness severity[74]. The zinc supplementation bunch likewise had a generous ascent in hair zinc levels in accordance with the past investigation, just as a critical improvement in the recurrence and seriousness of AD ($P=0.04$) comparative with the non-supplementation community[10]. One oral zinc supplementation randomized control preliminary (RCT) indicated an improvement in the degree and seriousness of AD among AD patients with zinc lack[75]. Our discoveries are viable with the past investigation that AD seriousness is related with low zinc [9].The consequences of these discoveries are predictable with different examinations revealed that Pb levels were fundamentally connected with seriousness of illness, low quality of life, and atopic dermatitis. The origin of lead admission is unsure that most food items are found in old structures and divider paints, as well. Exact food shirking movement was consequently remarkably typical among patients with AD[76, 77], while there was conflict with our information, the examination by Kim et al. (2019) found that AD seriousness was not fundamentally connected with Pb lead level[78].

Conclusion

- Through this investigation , it was discovered that the degree of schooling, position and family ancestry assume a critical function through skin inflammation pervasiveness.
- We found that the assessment of the zinc and lead serum levels in the dermatitis populace might be valuable, and further examination of their immunological capacity in pathogenesis and treatment of skin inflammation is required.
- Changes in patients' groupings of Mg, Ti, Mn, Fe , Zn, Al, Cd, Cr , Cu , Ni, Pb, and Se as stood out from the sound network demonstrated the chance of utilizing these components in giving treatment reactions.

References

- [1] Yong AM-Y, Tay Y-K. Atopic dermatitis: racial and ethnic differences. *Dermatologic clinics*. 2017;35:395-402.
- [2] Owens L, Laing IA, Zhang G, Turner S, Le Souëf PN. Prevalence of allergic sensitization, hay fever, eczema, and asthma in a longitudinal birth cohort. *Journal of asthma and allergy*. 2018;11:173.
- [3] Karabacak E, Aydin E, Kutlu A, Ozcan O, Muftuoglu T, Gunes A, et al. Erythrocyte zinc level in patients with atopic dermatitis and its relation to SCORAD index. *Advances in Dermatology and Allergology/Postępy Dermatologii i Alergologii*. 2016;33:349.
- [4] Strucinska M, Rowicka G, Riahi A. Profiles of selected nutrients affecting skin condition in children with atopic dermatitis. *Roczniki Państwowego Zakładu Higieny*. 2015;66.
- [5] Wada O. What are Trace Elements? 2004.
- [6] Bhattacharya PT, Misra SR, Hussain M. Nutritional aspects of essential trace elements in oral health and disease: an extensive review. *Scientifica*. 2016;2016.
- [7] A Puertollano M, Puertollano E, Alvarez de Cienfuegos G, A de Pablo M. Dietary antioxidants: immunity and host defense. *Current topics in medicinal chemistry*. 2011;11:1752-66.
- [8] Prashanth L, Kattapagari KK, Chitturi RT, Baddam VRR, Prasad LK. A review on role of essential trace elements in health and disease. *Journal of dr ntr university of health sciences*. 2015;4:75.
- [9] Vaughn AR, Foolad N, Maarouf M, Tran KA, Shi VY. Micronutrients in atopic dermatitis: a systematic review. *The Journal of Alternative and Complementary Medicine*. 2019;25:567-77.
- [10] Kim JE, Yoo SR, Jeong MG, Ko JY, Ro YS. Hair zinc levels and the efficacy of oral zinc supplementation in children with atopic dermatitis. *Acta dermato-venereologica*. 2014;94:558-62.
- [11] Toyran M, Kaymak M, Vezir E, Harmanci K, Kaya A, Giniş T, et al. 3 Trace Element Levels in Children With Atopic Dermatitis. *Journal of Investigational Allergology and Clinical Immunology*. 2012;22:341.
- [12] Ta V, Chin WK, White AA. Allergic contact dermatitis to testosterone and estrogen in transdermal therapeutic systems. *Dermatitis*. 2014;25:279.
- [13] Thyssen J, Johansen J, Linneberg A, Menné T, Engkilde K. The association between contact sensitization and atopic disease by linkage of a clinical database and a nationwide patient registry. *Allergy*. 2012;67:1157-64.
- [14] Jedrychowski W, Perera F, Maugeri U, Miller RL, Rembiasz M, Flak E, et al. Intrauterine exposure to lead may enhance sensitization to common inhalant allergens in early childhood: a prospective prebirth cohort study. *Environmental research*. 2011;111:119-24.

- [15] Eichenfield LF, Hanifin JM, Luger TA, Stevens SR, Pride HB. Consensus conference on pediatric atopic dermatitis. *Journal of the American Academy of Dermatology*. 2003;49:1088-95.
- [16] Kunz B, Oranje A, Labreze L, Stalder J-F, Ring J, Taieb A. Clinical validation and guidelines for the SCORAD index: consensus report of the European Task Force on Atopic Dermatitis. *Dermatology*. 1997;195:10-9.
- [17] Bonamonte D, Filoni A, Vestita M, Romita P, Foti C, Angelini G. The Role of the Environmental Risk Factors in the Pathogenesis and Clinical Outcome of Atopic Dermatitis. *BioMed research international*. 2019;2019.
- [18] Osman NO. Investigation of selected trace elements in hair samples of eczema patients. 2010.
- [19] Khan S, Shaikh M, Ali M. Depression in common dermatological patients attending outpatient department. *KYAMC Journal*. 2012;2:164-71.
- [20] Ahn K. The role of air pollutants in atopic dermatitis. *Journal of Allergy and Clinical Immunology*. 2014;134:993-9.
- [21] Eder W, Ege MJ, von Mutius E. The asthma epidemic. *New England Journal of Medicine*. 2006;355:2226-35.
- [22] Ebbehøj NE, Agner T, Zimerson E, Bruze M. Outbreak of eczema and rhinitis in a group of office workers in Greenland. *International journal of circumpolar health*. 2015;74:27919.
- [23] Chinn D, Poyner T, Sibley G. Randomized controlled trial of a single dermatology nurse consultation in primary care on the quality of life of children with atopic eczema. *British Journal of Dermatology*. 2002;146:432-9.
- [24] Staab D, Diepgen TL, Fartasch M, Kupfer J, Lob-Corzilius T, Ring J, et al. Age related, structured educational programmes for the management of atopic dermatitis in children and adolescents: multicentre, randomised controlled trial. *Bmj*. 2006;332:933-8.
- [25] Schram M, Tedja A, Spijker R, Bos J, Williams H, Spuls PI. Is there a rural/urban gradient in the prevalence of eczema? A systematic review. *British Journal of Dermatology*. 2010;162:964-73.
- [26] Flohr C, Pascoe D, Williams H. Atopic dermatitis and the 'hygiene hypothesis': too clean to be true? *British journal of dermatology*. 2005;152:202-16.
- [27] English J, Dawe R, Ferguson J. Environmental effects and skin disease. *British medical bulletin*. 2003;68:129-42.
- [28] Hugg T, Ruotsalainen R, Jaakkola MS, Pushkarev V, Jaakkola JJ. Comparison of allergic diseases, symptoms and respiratory infections between Finnish and Russian school children. *European journal of epidemiology*. 2008;23:123-33.
- [29] Saunes M, Øien T, Storror O, Johnsen R. Family eczema-history in 2-year olds with eczema; a prospective, population-based study. The PACT-study, Norway. *BMC dermatology*. 2011;11:11.
- [30] Wadonda-Kabondo N, Sterne J, Golding J, Kennedy C, Archer C, Dunnill M. Association of parental eczema, hayfever, and asthma with atopic dermatitis in infancy: birth cohort study. *Archives of disease in childhood*. 2004;89:917-21.
- [31] Chu H, Shin JU, Park CO, Lee H, Lee J, Lee KH. Clinical diversity of atopic dermatitis: a review of 5,000 patients at a single institute. *Allergy, asthma & immunology research*. 2017;9:158-68.
- [32] Wananukul S, Chatproedprai S, Tempark T, Phuthongkam W, Chatchatee P. The natural course of childhood atopic dermatitis: a retrospective cohort study. *Asian Pacific journal of allergy and immunology*. 2015;33.
- [33] Pyun BY. Natural history and risk factors of atopic dermatitis in children. *Allergy, asthma & immunology research*. 2015;7:101-5.

- [34] Kulthanan K, Samutrapong P, Jiamton S, Tuchinda P. Adult-onset atopic dermatitis: a cross-sectional study of natural history and clinical manifestation. *Asian Pacific journal of allergy and immunology*. 2007;25:207.
- [35] Tam M, Gomez S, Gonzalez-Gross M, Marcos A. Possible roles of magnesium on the immune system. *European journal of clinical nutrition*. 2003;57:1193-7.
- [36] Makiura M, Akamatsu H, Akita H, Yagami A, Shimizu Y, Eiro H, et al. Atopic dermatitis-like symptoms in HR-1 hairless mice fed a diet low in magnesium and zinc. *Journal of international medical research*. 2004;32:392-9.
- [37] Tuchinda P, Liu Y, Tammara A, Harberts E, Goldner R, Gaspari AA. Resolution of occupational dermatitis related to manganese exposures. *Dermatitis: contact, atopic, occupational, drug*. 2014;25:280-1.
- [38] Kim JE, Shin JM, Ko JY, Ro YS. Hair mineral analysis in children with atopic dermatitis. *Annals of dermatology*. 2017;29:251-3.
- [39] AFRIDI HI, Kazi TG, Jamali MK, KAZI GH, Shar GQ. The status of trace and toxic elements in biological samples (scalp hair) of skin-disease patients and normal subjects. *Turkish Journal of Medical Sciences*. 2006;36:223-30.
- [40] Kaur S, Eisen M, Zilmer M, Kullisaar T, Rehema A, Vihalemm T. Patients with allergic and irritant contact dermatitis are characterized by striking change of iron and oxidized glutathione status in nonlesional area of the skin. *Journal of investigative dermatology*. 2001;116:886-90.
- [41] Meding B, Lantto R, Lindahl G, Wrangsjö K, Bengtsson B. Occupational skin disease in Sweden—a 12-year follow-up. *Contact dermatitis*. 2005;53:308-13.
- [42] Lee SH, Huang JW, Hung KY, Leu LJ, Kan YT, Yang CS, et al. Trace Metals' abnormalities in hemodialysis patients: relationship with medications. *Artificial Organs*. 2000;24:841-4.
- [43] Aquino M, Mucci T. Systemic contact dermatitis and allergy to biomedical devices. *Current allergy and asthma reports*. 2013;13:518-27.
- [44] Malajian D, Belsito DV. Cutaneous delayed-type hypersensitivity in patients with atopic dermatitis. *Journal of the American Academy of Dermatology*. 2013;69:232-7.
- [45] Thyssen JP, Skare L, Lundgren L, Menné T, Johansen JD, Maibach HI, et al. Sensitivity and specificity of the nickel spot (dimethylglyoxime) test. *Contact dermatitis*. 2010;62:279-88.
- [46] Jensen P, Thyssen JP, Johansen JD, Skare L, Menné T, Lidén C. Occupational hand eczema caused by nickel and evaluated by quantitative exposure assessment. *Contact dermatitis*. 2011;64:32-6.
- [47] David T, Wells F, Sharpe T, Gibbs A, Devlin J. Serum levels of trace metals in children with atopic eczema. *British Journal of Dermatology*. 1990;122:485-9.
- [48] Tasaki M, Hanada K, Hashimoto I. Analyses of serum copper and zinc levels and copper/zinc ratios in skin diseases. *The Journal of dermatology*. 1993;20:21-4.
- [49] ERTUNÇ V, PARLAK M, Şenol D, AKTAŞ A, KARAKUZU A, ALTUNTAŞ İ. The serum levels of some important elements and alkaline phosphatase in patients with atopic dermatitis. *TURKISH JOURNAL OF MEDICAL SCIENCES*. 1996;26:593-6.
- [50] El-Kholy M, Gas MA, El-Shimi S, El-Baz F, El-Tayeb H, Abdel-Hamid M. Zinc and copper status in children with bronchial asthma and atopic dermatitis. *The Journal of the Egyptian Public Health Association*. 1990;65:657-68.
- [51] Chen C-J, Lai J-S, Wu C-C, Lin T-S. Serum selenium in adult Taiwanese. *Science of the total environment*. 2006;367:448-50.

- [52] Benes B, Spevackova V, Smid J, Batariova A, Cejchanova M, Zitkova L. Effects of age, BMI, smoking and contraception on levels of Cu, Se and Zn in the blood of the population in the Czech Republic. *Central European journal of public health*. 2005;13:202.
- [53] Clark NA, Teschke K, Rideout K, Copes R. Trace element levels in adults from the west coast of Canada and associations with age, gender, diet, activities, and levels of other trace elements. *Chemosphere*. 2007;70:155-64.
- [54] Ghayour-Mobarhan M, Taylor A, New S, Lamb D, Ferns G. Determinants of serum copper, zinc and selenium in healthy subjects. *Annals of clinical biochemistry*. 2005;42:364-75.
- [55] Gammoh NZ, Rink L. Zinc in infection and inflammation. *Nutrients*. 2017;9:624.
- [56] Metryka E, Chibowska K, Gutowska I, Falkowska A, Kupnicka P, Barczak K, et al. Lead (Pb) exposure enhances expression of factors associated with inflammation. *International journal of molecular sciences*. 2018;19:1813.
- [57] Rocha G, Steinbach C, Munhoz J, Madia M, Faria J, Hoeltgebaum D, et al. Trace metal levels in serum and urine of a population in southern Brazil. *Journal of Trace Elements in Medicine and Biology*. 2016;35:61-5.
- [58] Kim H-J, Lim H-S, Lee K-R, Choi M-H, Kang NM, Lee CH, et al. Determination of trace metal levels in the general population of Korea. *International journal of environmental research and public health*. 2017;14:702.
- [59] Hotz C, Peerson JM, Brown KH. Suggested lower cutoffs of serum zinc concentrations for assessing zinc status: reanalysis of the second National Health and Nutrition Examination Survey data (1976–1980). *The American journal of clinical nutrition*. 2003;78:756-64.
- [60] Harding KL, Aguayo VM, Masters WA, Webb P. Education and micronutrient deficiencies: an ecological study exploring interactions between women’s schooling and children’s micronutrient status. *BMC public health*. 2018;18:470.
- [61] Lee JW, Lee CK, Moon CS, Choi IJ, Lee KJ, Yi S-M, et al. Korea National Survey for Environmental Pollutants in the Human Body 2008: heavy metals in the blood or urine of the Korean population. *International journal of hygiene and environmental health*. 2012;215:449-57.
- [62] Son J-Y, Lee J, Paek D, Lee J-T. Blood levels of lead, cadmium, and mercury in the Korean population: results from the Second Korean National Human Exposure and Bio-monitoring Examination. *Environmental research*. 2009;109:738-44.
- [63] Yaylalı-Abanuz G. Heavy metal contamination of surface soil around Gebze industrial area, Turkey. *Microchemical Journal*. 2011;99:82-92.
- [64] Mohmand J, Eqani SAMAS, Fasola M, Alamdar A, Mustafa I, Ali N, et al. Human exposure to toxic metals via contaminated dust: Bio-accumulation trends and their potential risk estimation. *Chemosphere*. 2015;132:142-51.
- [65] Kiec-Swierczynska M, Krecisz B, Chomiczewska D, Sobala W. Trends in allergy to the 10 most frequent contact allergens in patients examined at the Nofer Institute, Lodz, Poland in 1996-2009. *Postępy Dermatologii i Alergologii*. 2012;29:19.
- [66] Cabral Pinto M, Marinho-Reis P, Almeida A, Pinto E, Neves O, Inácio M, et al. Links between Cognitive Status and Trace Element Levels in Hair for an Environmentally Exposed Population: A Case Study in the Surroundings of the Estarreja Industrial Area. *International journal of environmental research and public health*. 2019;16:4560.

- [67] Yamada T, Saunders T, Kuroda S, Sera K, Nakamura T, Takatsuji T, et al. Cohort study for prevention of atopic dermatitis using hair mineral contents. *Journal of Trace Elements in Medicine and Biology*. 2013;27:126-31.
- [68] Hashemipour M, Kelishadi R, Shapouri J, Sarrafzadegan N, Amini M, Tavakoli N, et al. Effect of zinc supplementation on insulin resistance and components of the metabolic syndrome in prepubertal obese children. *Hormones*. 2009;8:279-85.
- [69] Jing M-Y, Sun J-Y, Weng X-Y. Insights on zinc regulation of food intake and macronutrient selection. *Biological trace element research*. 2007;115:187-94.
- [70] Baltaci AK, Mogulkoc R. Leptin and zinc relation: in regulation of food intake and immunity. *Indian journal of endocrinology and metabolism*. 2012;16:S611.
- [71] An HC, Sung JH, Lee J, Sim CS, Kim SH, Kim Y. The association between cadmium and lead exposure and blood pressure among workers of a smelting industry: a cross-sectional study. *Annals of occupational and environmental medicine*. 2017;29:47.
- [72] Skalnaya MG, Tinkov AA, Demidov VA, Serebryansky EP, Nikonorov AA, Skalny AV. Hair toxic element content in adult men and women in relation to body mass index. *Biological trace element research*. 2014;161:13-9.
- [73] Skolarczyk J, Pekar J, Łabędź D, Skórzyńska-Dziduszko K. Role of heavy metals in the development of obesity: a review of research. *Journal of Elementology*. 2018;23.
- [74] Hon KLE, Wang SS, Hung EC, Lam HS, Lui HH, Chow CM, et al. Serum levels of heavy metals in childhood eczema and skin diseases: friends or foes. *Pediatric allergy and immunology*. 2010;21:831-6.
- [75] Gray N, Dhana A, Stein D, Khumalo N. Zinc and atopic dermatitis: a systematic review and meta-analysis. *Journal of the European Academy of Dermatology and Venereology*. 2019;33:1042-50.
- [76] Hon K, Leung T, Kam W, Lam M, Fok T, Ng P. Dietary restriction and supplementation in children with atopic eczema. *Clinical and Experimental Dermatology: Clinical dermatology*. 2006;31:187-91.
- [77] Hon KIE, Ma KC, Wong Y, Leung TF, Fok TF. A survey of traditional Chinese medicine use in children with atopic dermatitis attending a paediatric dermatology clinic. *Journal of dermatological treatment*. 2005;16:154-7.
- [78] Kim J, Kim S, Woo S-y, Chung J-Y, Hong Y-S, Oh S-Y, et al. Prenatal Exposure to Lead and Chromium is Associated with IL-13 Levels in Umbilical Cord Blood and Severity of Atopic Dermatitis: COCOA Study. *Immune Network*. 2019;19.